

ACCRETION ONTO NEARBY SUPERMASSIVE BLACK HOLES: *CHANDRA* CONSTRAINTS ON THE DOMINANT CLUSTER GALAXY NGC 6166

TIZIANA DI MATTEO¹

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138; tdimatteo@cfa.harvard.edu

RODERICK M. JOHNSTONE, STEVEN W. ALLEN AND ANDREW C. FABIAN

Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK; rmj,swa,acf@ast.cam.ac.uk

Draft version February 1, 2008

ABSTRACT

Chandra observations of low-luminosity supermassive black holes in nearby elliptical galaxies provide tight limits on both their nuclear luminosities and on their Bondi accretion rates. We examine *Chandra* constraints on NGC 6166, the dominant galaxy in the cluster Abell 2199, which hosts a $\sim 10^9 M_\odot$ black hole. We measure a nuclear X-ray luminosity $L_{x,1\text{ keV}} \sim 10^{40} \text{ erg s}^{-1}$ and show that the density and temperature profiles of the hot interstellar medium imply a Bondi accretion rate of $\dot{M}_{\text{Bondi}} \lesssim 3 \times 10^{-2} M_\odot \text{ yr}^{-1}$. This accretion rate predicts a nuclear luminosity of $\sim 10^{44} \text{ erg s}^{-1}$ for a canonical radiative efficiency of 10%. Unless the Bondi estimate is inappropriate and/or the accretion rate onto the black hole is significantly reduced, the observed nuclear flux constrains the radiative efficiency of the accretion flow to be $\eta \approx 10^{-5}$. We show that low-radiative efficiency accretion flows with radial density profiles $\rho \propto r^{-3/2}$ (and not significantly flatter) can explain the observed nuclear X-ray luminosity but that the power output from the jets in NGC 6166 is also important to the energetics of the system.

1. INTRODUCTION

Recent dynamical measurements of black hole masses in nearby galaxies have transformed black hole studies, allowing us to quantify directly their expected accretion power. In those cases in which the fuel supply to the black holes can be estimated, it has become apparent that the expected power far exceeds the observed luminosity. This situation is perhaps best illustrated by the case of nearby elliptical galaxy nuclei (e.g. Fabian & Rees 1995; Di Matteo et al. 2000a, hereafter DM00a and references therein) and our Galactic center (e.g. Narayan et al. 1998 and references therein). There are only two possible explanations for the low-luminosities of nearby black holes: (a) the accretion occurs at extremely low rates or (b) the accretion occurs at low radiative efficiencies as predicted, for example, by advection dominated accretion models (e.g. Rees et al. 1982; Narayan & Yi 1994, 1995; Abramowicz et al. 1995).

Assuming that the accretion in elliptical galaxies occurs primarily from the hot, quasi-spherical interstellar medium (ISM), Bondi theory (1952) can be used to estimate the accretion rates onto the supermassive black holes. Such estimates require accurate measurements of both the density and temperature of the hot ISM at the 'Bondi accretion radius', the radius where the gravitational potential of the central black holes begins to dominate the dynamics of the hot gas.

In order to unambiguously determine whether the low luminosities of nearby black holes are due to a low radiative efficiency in the accreting gas, it is also necessary to measure the nuclear power. When combined with the estimated accretion rates, this gives us a direct measurement of the radiative efficiency of accretion, η .

Thanks to its high spatial resolution and sensitivity, the

Chandra X-ray observatory is able, for the first time, to detect nuclear X-ray point sources in nearby galaxies and provide us with direct measurements of their luminosities. *Chandra* also allows us to measure the central densities and temperatures of the ISM close to the accretion radii of the central black holes, and therefore determine the Bondi accretion rates in these systems to much greater accuracy than before.

In this *Letter* we explore the implications of *Chandra* observations of the giant elliptical galaxy NGC 6166. We report the detection of a nuclear point source in the galaxy with a luminosity $L_x \sim$ a few $10^{40} \text{ erg s}^{-1}$. We measure the central X-ray gas density and temperature and calculate the Bondi accretion rate. We show that if the central black hole of $\sim 10^9 M_\odot$ is fed at the estimated Bondi rate, the inferred accretion radiative efficiency is $\eta \lesssim 10^{-5}$. We show that at the given accretion rate ADAF models can explain the observed nuclear luminosity. The presence of outflows in the accretion flows is also consistent with the present constraints. We discuss possible reasons why the Bondi accretion rate estimate might be inappropriate and show that a significant contribution to the nuclear flux is likely to be due to emission from the jets.

2. NUCLEAR LUMINOSITY AND ACCRETION RATE

NGC 6166 ($z = 0.031$) is a cD galaxy at the center of the cooling flow cluster A2199, and is therefore located in a highly gas-rich environment. The central black hole mass is estimated by Ferrarese & Merritt (2000) using the correlation between the masses of black holes and the velocity dispersions of their host bulges ($M_{\text{BH}} \propto \sigma_c^{4.7}$). For NGC 6166, $\sigma_c = 326 \text{ km s}^{-1}$ which gives $M_{\text{BH}} \sim 10^9 M_\odot$, implying a discrepancy of about an order of magnitude with

¹ *Chandra* Fellow

the value obtained by Magorrian et al. (1998), who estimated $M_{BH} \sim 10^{10} M_{\odot}$.

NGC 6166 is known to host an active nucleus, classified as an FR I source, which powers two symmetric parsec-scale radio jets and radio lobes. The jets are close to the plane of the sky (Giovannini et al. 1998). Beaming should not therefore be important. Given the measured black hole mass, the Eddington luminosity of NGC 6166 is $L_{\text{Edd}} \sim 10^{47} \text{ erg s}^{-1}$. However, the observed bolometric luminosity is $\lesssim 10^{41} \text{ erg s}^{-1}$.

We adopt the cosmological parameters $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega = 1$ and $\Lambda = 0$, which imply a distance to NGC 6166 of 142 Mpc).

2.1. L_x

The *Chandra* observations of the A2199 cluster were carried out on 1999 December 11 using the Advanced CCD Imaging Spectrometer and back-illuminated CCD detectors. The net exposure time was 14.95 ks. Here, we only highlight results for the central, dominant galaxy NGC 6166; details of the cluster and point source analysis are discussed by Johnstone et al. (2000).

The *Chandra* data for NGC 6166 reveal the presence of a nuclear point-like X-ray source (detected over background at 7 sigma significance) with a surface brightness distribution consistent with the point-spread function (PSF) of the instrument ($\sigma \sim 0.45 \text{ arcsec}$). The X-ray source is coincident with the peak of the emission from the central radio source (Giovannini et al. 1998). In order to accurately determine the central point source flux, we have modeled the central surface brightness profile using a simple power-law ($I_X \propto r^{-\alpha}$; accounting for the underlying ISM) plus Gaussian model, convolved with the instrumental PSF. We find that the surface brightness profile within the innermost 10 kpc radius region can be well-described by a power-law model of slope $\alpha = 0.4 \pm 0.025$ and a point source with 55 ± 27 counts (1σ errors; where the error in the counts is dominated by the uncertainty in the slope of the density profile).

We have extracted a spectrum for the central source using a region of $\sim 1.5 \text{ arcsec}$ radius (which should contain $\gtrsim 90$ per cent of the flux from the source) and used a background spectrum taken from the 1.5 – 2.5 arcsec annulus. Given the faintness of the central source and the moderate exposure time of the observation, the spectral parameters are not firmly constrained and the data can be described by either a power law or a thermal plasma model (mekal). The power-law model gives a photon index $\Gamma = 1.54_{-0.45}^{+0.46}$ assuming Galactic absorption, and a flux density at 1 keV of $7 \pm 2 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$, where all parameters are quoted at a 1σ confidence. (Allowing for intrinsic absorption, the maximum intrinsic column density is $N_H \sim 2 \times 10^{21} \text{ atom cm}^{-2}$ for $\Gamma = 2.9$ and $F(1 \text{ keV}) = 14 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$; see Fig. 3 and contour plots in Johnstone et al. 2000). The flux determined from the spectrum is consistent with the value measured from the surface brightness profile, assuming a consistent spectral model for the nuclear emission.

2.2. $\dot{M}$

We can estimate the accretion rate from the hot ISM onto the central black hole using Bondi accretion the-

ory. Matter passing within the accretion radius $R_A \approx (c/c_s)^2 R_S$ is assumed to be accreted. Here c is the speed of light, $c_s \sim 10^4 T^{1/2} \text{ cm s}^{-1}$ is the sound speed (with T the gas temperature), and $R_S = 2GM/c^2$ the Schwarzschild radius of the black hole.

We have measured the temperature and density profile of the intracluster gas within the central 100 kpc radius region. Direct measurements of these quantities are only possible down to within $r \sim 1 \text{ arcsec}$ of the nucleus, due to contamination from the nuclear point source. The filled triangles in Fig. 1 show the temperature profile determined by fitting the annular counts with a single temperature thermal emission model (mekal) acted on by Galactic and (free-fitting) intrinsic absorption. The open squares show the deprojected temperature profile in which the inner regions are corrected for the emission from further out seen in projection. The temperature is higher in the outer regions, where the cluster potential dominates and decreases (and flattens off) in the inner regions where the galaxy potential takes over.

The density profile shown in Fig. 2 was derived by deprojecting the surface brightness profile (e.g. Fabian et al. 1981) and matching the deprojected temperature profile. The density profile (Fig. 2) is steeper in the outer regions ($r \gtrsim 10 \text{ kpc}$) and follows the canonical cooling-flow profile of $\rho \propto r^{-1}$. Inside this radius the density profile flattens to a value $\rho \propto r^{-0.45 \pm 0.09}$ (we exclude the central 1.5 arcsec region, which is affected by the point source).

We take $kT \sim 1.3 \text{ keV}$ as measured in the innermost regions. This gives

$$R_A \sim 0.06 T_{1.3}^{-1} M_9 \text{ kpc}, \quad (1)$$

where $M_9 = 10^9 M_{\odot}$ which corresponds to $\sim 0.08 \text{ arcsec}$.

The accretion rate is related to the density and temperature by the continuity equation, with the velocity $v = c_s$;

$$\dot{M} = 4\pi R_A^2 \rho_A c_s(R_A) \quad (2)$$

where ρ_A and $c_s(R_A)$ are the density and sound speed at the accretion radius.

Given that the density and temperature cannot be measured directly at the accretion radius we assume $kT = 1.3_{-0.7}^{+0.6} \text{ keV}$ and estimate the density at R_A by extrapolating the best-fit power-law model (see above) which gives $n(R_A) \sim 0.6 \pm 0.1 \text{ cm}^{-3}$.

This implies

$$\begin{aligned} \dot{M}_{\text{Bondi}} &= 1.3_{-0.7}^{+1.2} \times 10^{24} M_9^2 T_{1.3}^{-3/2} n_{0.6} \text{ g s}^{-1} \quad (3) \\ &\sim 3 \times 10^{-2} M_{\odot} \text{ yr}^{-1}, \end{aligned}$$

(where we have neglected the uncertainties on M_{BH}). Eqn. (2) gives an estimate of the accretion at the outer edge of the accretion flow. As discussed in the next session, the mass accretion rate onto the black hole may be smaller if e.g. it decreases with radius because of an out-flow.

This Bondi accretion rate implies $L_{\text{Bondi}} = \eta \dot{M}_{\text{Bondi}} c^2 \sim 10^{44} \text{ erg s}^{-1}$ if $\eta = 0.1$, as in a standard radiatively efficient thin disk. This is about four orders of magnitude greater than the observed luminosity of the central X-ray source, implying that the radiative efficiency has to be $\eta \sim 10^{-5}$ suggesting that hot accretion flows with low radiative efficiencies may be relevant.

3. MODELS

In a hot accretion flow around a supermassive black hole, the majority of the emission arises in the radio and X-ray bands. In the radio band the emission results from synchrotron radiation from the inner parts of the accretion flow. The X-ray emission is due to either bremsstrahlung emission or inverse Compton scattering of the soft synchrotron photons (e.g. Narayan, Barret & McClintock 1998).

We measure radii in the flow in Schwarzschild units: $R = rR_S$, where $R_S = 2GM_{BH}/c^2$. We measure black hole masses in solar units and accretion rates in Eddington units: $M_{BH} = m M_\odot$ and $\dot{M} = \dot{m} \dot{M}_{Edd}$. We take $\dot{M}_{Edd} = 10L_{Edd}/c^2 = 2.2 \times 10^{-8} m M_\odot \text{ yr}^{-1}$, i.e., with a canonical 10% efficiency. We take $r = 10^4$ to be the outer radius of the flow. The Bondi accretion rate in Eddington units is $\dot{m}_{Bondi} = 1.3 \times 10^{-3}$.

The predicted spectrum from an ADAF depends (weakly) on the ratio of gas to magnetic pressure β , the viscosity parameter α , and the fraction of the turbulent energy in the plasma which heats the electrons, δ . Here, we fix $\alpha = 0.1$, $\beta = 10$, and $\delta = 0.1$. The two major parameters, though, are the accretion rate $\dot{M}$ and the black hole mass M_{BH} , both of which are constrained. With M_{BH} assumed, we normalize the models to the observed *Chandra* flux. This gives us the $\dot{m}$ required by the models to explain the X-ray emission. This has to be consistent with our estimate of $\dot{m}_{Bondi}$.

The solid line in Fig. 3 shows the predicted spectrum for a pure inflow ADAF model fitted to the 1 keV flux. In this model $\dot{m} \sim 10^{-3}$, consistent with the Bondi estimate. Comptonization of the synchrotron emission dominates the X-ray emission in this model. ADAFs models can therefore explain the observed X-ray emission. At the given Bondi rate the radiative efficiency of an ADAF is $\sim 10^{-5}$ corresponding to that derived in §2.2. Note, however, that in these models the synchrotron component cannot explain the VLBI radio emission (Giovannini et al. 1998; Fig. 3). This implies that the majority of the contribution in the radio band is likely to be due to emission from the jets (e.g., dotted line).

Following the proposal of Blandford & Begelman (1999) that mass loss may be important in hot accretion flows, Di Matteo et al. (1999, hereafter DM99; DM00a) and Quataert & Narayan (1999; QN99) have constructed models in which a significant fraction of the accreted mass in an ADAF is lost to an outflow rather than being accreted onto the central object. In that work (see also Di Matteo, Fabian & Carilli 2000b, hereafter DM00b) it was shown that such models are favored (or required) over the pure inflow ADAF for explaining the observed spectral energy distributions of elliptical galaxy nuclei.

The importance of outflows in ADAFs can be parameterized by using different radial density profiles in the flows with $\rho \propto r^{-3/2+p}$ and $0 < p \leq 1$ (i.e. flatter than in the pure Bondi inflow with $\rho \propto r^{-3/2}$). This implies mass inflow rates which satisfy the relation $\dot{M} = \dot{M}_{out}(r/r_{out})^p$, where we should satisfy $\dot{M}_{out} \sim \dot{M}_{Bondi}$ and $r_{out} = 10^4$ as before. Bremsstrahlung emission in these models dom-

inated the X-ray emission. This produces a harder X-ray spectrum, most of which (in the 0.5 - 10 keV band) is produced in the outer regions of the flow (the temperatures in the inner regions approach 100 keV to 1 MeV at $r \sim r_{out}$ where $\dot{m}_{out} \sim \dot{m}$ (see details in DM00a; QN99). The dashed line shows a model with $p = 0.1$ which implies $\dot{m}_{out} = 3 \times 10^{-3}$. If $p \gtrsim 0.1$, $\dot{m}_{out}$ becomes inconsistent with the Bondi estimate (Eq.3) unless the errors on M_{BH} are sufficiently large (e.g., for $p = 0.2$, $\dot{m}_{out} \sim 6 \times 10^{-3}$, so M_{BH} would need to have been underestimated by a factor $\gtrsim 50\%$).

For certain regimes (e.g. for $\alpha \lesssim 0.1$) convection might also become important in hot accretion flows, (e.g. Stone, Pringle & Begelman 1999; Quataert & Gruzinov 2000; Narayan, Igumenshchev & Abramowicz 2000). Although CDAFs have a very different structure than ADAFs, they have density profiles $\rho \propto r^{-1/2}$ (corresponding to the $p = 1$ case of the outflow models above). Ball, Narayan & Quataert (2000) have constructed spectral models for CDAFs similar to those considered by DM99 and QN99 for outflows². Taking $p = 1$, even with $\dot{m}_{out} \sim \dot{m}_{Bondi}$, implies very small contributions from a CDAF to the X-ray flux. A CDAF if present cannot explain the observed emission.

Finally, the dotted line in Fig. 3 shows an example of non-thermal jet models for the full band spectrum. It is clear that the observed radio jets must contribute significantly to the observed radio and optical emission even if an ADAF (with or without outflows) is also present. Fig. 3 shows that the full spectral energy distribution of NGC 6166 can be easily explained by the emission from different compact regions emitting self-absorbed non-thermal synchrotron radiation. The typical parameters for these components are $B \sim 7 \times 10^{-4} \nu_{tGHz} T_{B12}^{-2} \text{ G}$ and $\tau_T \sim 0.4 T_{B12}^5 \nu_{tGHz}$; where $T_{B12} = 10^{12} \text{ K}$ is the brightness temperature and ν_t the self-absorption frequency (see Di Matteo, Carilli & Fabian 2000 for details, hereafter DM00b). Emission from a jet is far more efficient even if it originates from relatively low density regions and can easily give rise to the observed X-ray flux (see also e.g. Falcke 1999). Explaining most of the observed emission as being due to the jets would be consistent with the unifying model in which FR I radio galaxies are mis-oriented BL Lac objects and the fact that the radio-to-optical and optical-to-X-ray spectral indexes of NGC 6166 are consistent with those of radio selected BL Lac objects (see Fig. 3 in Hardcastle & Worrall 1999). However, this interpretation would still not relax the requirement for a low-radiative efficiency in the accretion flow.

4. DISCUSSION

We have examined *Chandra* constraints on the low luminosity active nucleus in NGC 6166. By adopting $M_{BH} = 10^9 M_\odot$ and inferring the ISM density and temperature at R_A we have estimated the Bondi accretion rate to be $\sim 3 \times 10^{-2} M_\odot \text{ yr}^{-1}$. This value is $\lesssim 10$ times lower than the typical Bondi rates determined in previous studies based on ROSAT High Resolution Imager (HRI) observations of other nearby elliptical nuclei (e.g., M87, NGC4649, NGC4472; DM00a) and is more consistent with

²Note that in a CDAF the Bondi accretion rate estimate is not appropriate as convection and possibly conduction significantly alter the rate at which matter is fed into the flow. The accretion rates (at large radii) are expected to be lower than in the ADAFs

the upper limits inferred from high frequency radio studies of the accretion rates at small radii in the ellipticals (DM99;DM00b; Wrobel & Herrnstein 2000).

The *Chandra* observations of NGC 6166 indicate a nuclear luminosity of $L_{x,0.5-7\text{ keV}} \sim 4 \times 10^{40} \text{ erg s}^{-1}$. This implies that if matter is accreted from the ISM at the Bondi rate the radiative efficiency of the accretion flow has to be $\eta \lesssim 10^{-5}$. We have shown that this is consistent with the predictions of pure inflow ADAF models. Hot accretion flow models with significantly flatter density profiles (with strong outflows or CDAFs), which are favored on theoretical grounds and from radio observations (DM99,DM00b), cannot explain the observed nuclear flux (unless M_{BH} is significantly larger). This may imply that most of the observed flux is dominated by non-thermal emission from the jets.

Although with *Chandra* observations we can infer Bondi accretion rates with much greater confidence than before, for NGC 6166 we can only measure the ISM densities and temperatures directly at a radius $\sim 1 \text{ kpc}$ which is still a factor ~ 15 larger than R_A (§2.2). Given the distance to NGC 6166, previous ROSAT HRI observations would have only been able to measure the density profile down to $r \gtrsim 10 \text{ kpc}$, where we observe the density profile to steepen. Extrapolation of this steeper density profile to the accretion radius would then have lead to higher inferred accretion rates. Even with *Chandra*, however, we can still only infer the ISM density at R_A in NGC 6166, which implies that we could still be overestimating the Bondi accretion rate (if e.g. the density profile flattened further towards R_A). Given that in order to explain

the observed nuclear flux, an ADAF requires accretion rates similar to those currently estimated, a decrease in the Bondi estimate would make the presence of an ADAF undetectable and their emission negligible. Similar arguments apply if the temperature closer to R_A increased.

It is also possible that the Bondi calculation itself may be inappropriate. If strong convection were important, as in CDAFs, the rates at which matter is fed through the accretion radius could be significantly reduced. Heating of the ISM by the jets (DM00b) could also decrease significantly the Bondi accretion radius and hence the rate at which mass is fed into the accretion flows. If the accretion rate is low enough, the requirement for low radiative efficiency in the accretion flow may be significantly relaxed.

We have shown that emission from the jets in these low-luminosity nuclei provides an important contribution to the overall energy output of NGC 6166. Determining the variability properties of the X-ray emission may be important for resolving the jet contribution from that of an ADAF. *Chandra* observations of other, closer, low-luminosity black holes (e.g., M87) are needed to further assess the question of low radiative-efficiency versus low- $\dot{M}$ accretion in nearby black holes.

We thank Ramesh Narayan for many useful comments. T.D.M. acknowledges support for this work provided by NASA through Chandra Postdoctoral Fellowship grant number PF8-10005 awarded by the Chandra Science Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-39073. SWA and ACF thank the Royal Society for support.

REFERENCES

- Abramowicz M., Chen X., Kato S., Lasota J. P., Regev O., 1995, *ApJ*, 438, L37
 Ball G.H., Narayan R., Quataert E., 2000, *Apj*, submitted (astro-ph/0007037)
 Blandford R.D., Begelman M.C., 1999, *MNRAS*, 303, L1
 Chiaberge M., Capetti A., Celotti A., 1999, *A&A*, 349, 77
 Di Matteo T., Carilli C.L., Fabian A.C., 2000, *ApJ*, in press (DM00b)
 Di Matteo T., Fabian A.C., Rees M.J., Carilli C., Ivison R J., 1999, *MNRAS*, 305, 49
 Di Matteo, T., Quataert, E., Allen, S.W., Narayan, R., Fabian, A. C., 2000, *MNRAS*, 311, 507 (DM00a)
 Fabian A. C., Hu E.M., Cowie L.L., Grindlay J., 1981, *ApJ*, 248, 47
 Fabian A. C., Canizares C. R., 1988, *Nature*, 333, 829
 Falcke, H., 1999, in *The Central Parsecs of the Galaxy*, eds. H. Falcke et al. ASP Conf. Series, Vol. 186, p.148 (astro-ph/9909439)
 Ferrarese, L., Merritt, D., 2000, *ApJ*, 539, L9
 Giovannini G., Cotton W.D., Feretti L., Lara L., Venturi T., 1998, *ApJ*, 493, 632
 Hardcastle H.J., Worrall D.M., 1999, *MNRAS*, 309, 969
 Igumenshchev, I. V. Abramowicz, M. A. 1999, *MNRAS*, 303, 309
 Johnstone et al. 2000, in preparation
 Magorrian, J. et al. 1998, *AJ*, 115, 2285
 Mahadevan R., 1997, *ApJ*, 477, 585
 Narayan R., Yi I., 1995, *ApJ*, 444, 231
 Narayan R., Igumenshchev, I. V. Abramowicz, M. A., 2000, *ApJ*, 539, 798
 Narayan, R., Mahadevan, R. & Quataert, E. 1998, in *Theory of Black Hole Accretion Disks*, Eds. M. Abramowicz, G. Bjornsson, and J. Pringle (Cambridge: Cambridge University Press), p148
 Quataert E., Gruzinov I., 2000, *ApJ*, 539, 809
 Quataert E., Narayan R., 1999, *ApJ*, 520, 298
 Rees M. J., Begelman M. C., Blandford R. D., Phinney E. S., 1982, *Nature*, 295, 17
 Stone, J. M., Pringle, J. E. Begelman, M. C. 1999, *MNRAS*, 310, 1002
 Wrobel, J. M., Herrnstein, J. R, 2000, *ApJ*, 533, L111




Fig. 1.— The temperature profile in NGC 6166 measured by *Chandra*. The open squares show the deprojected temperature measurements and the filled triangles the observed (projected) values.




Fig. 2.— The deprojected density profile in the inner regions of A 2199.




Fig. 3.— Spectral models calculated for hot accretion flows fitted to the *Chandra* X-ray flux. The solid line is an ADAF model. The required accretion rate, $\dot{m} = 8 \times 10^{-4}$, is consistent with the inferred Bondi value. The dashed line shows a model with outflows. Here $p = 0.1$, $\dot{m}_{\text{out}} = 3 \times 10^{-3}$. The dotted line is the sum of self-absorbed, non-thermal synchrotron emitting regions in the jets. Beaming is ignored since the jets lie very close to the plane of the Sky (Giovannini et al. 1998). The solid dots are VLBI flux measurements from Giovannini et al. (1998), the open circles the VLBI core peak fluxes and the HST (Chiaberghe et al. 1999) continuum flux and ROSAT HRI upper limit (Hardcastle & Worrall 1999). The bow-tie shows the *Chandra* flux (in the 0.5 – 7 keV band) for the range of power-law slopes with the hatched area corresponding to the ones acceptable with Galactic absorption fixed.